

TITLE

A comparison of methods to estimate anaerobic capacity: Accumulated oxygen deficit and W' during constant and all-out work-rate profiles

AUTHOR

Muniz-Pumares, Daniel; Pedlar, Charles; Godfrey, R.; et al.

JOURNAL

Journal of Sports Sciences

DATE DEPOSITED

6 September 2017

This version available at

<https://research.stmarys.ac.uk/id/eprint/1748/>

COPYRIGHT AND REUSE

Open Research Archive makes this work available, in accordance with publisher policies, for research purposes.

VERSIONS

The version presented here may differ from the published version. For citation purposes, please consult the published version for pagination, volume/issue and date of publication.

Title: A comparison between methods to estimate anaerobic capacity: Effects of pacing on the accumulated oxygen deficit and the power-duration relationship

Running title: AOD and W' during constant-load and all-out exercise

Authors: Muniz-Pumares, Daniel^{1,2}; Pedlar, Charles¹; Godfrey, Richard³; Glaister, Mark¹

¹ School of Sport, Health and Applied Science, St Mary's University, Twickenham, UK.

² School of Life and Medical Sciences, University of Hertfordshire, Hatfield, UK.

³ The Centre for Sports Medicine and Human Performance, Brunel University, Uxbridge, UK.

Corresponding author: Daniel Muniz-Pumares

Department of Psychology and Sport Science

School Life and Medical Sciences

College Lane

University of Hertfordshire

Hatfield

AL10 9EU

United Kingdom

Telephone: [0170 728 3495](tel:01707283495)

Email: d.muniz@herts.ac.uk

Abstract

This study investigated i) whether the accumulated oxygen deficit (AOD) and the curvature constant of the power-duration relationship (W') remain unchanged during constant work-rate to exhaustion (CWR) and 3-min all-out (3MT) tests; and ii) the relationship between AOD and W' during CWR and 3MT. Twenty-one male cyclists (age: 40 ± 6 years; maximal oxygen uptake ($\dot{V}O_{2\max}$): 58 ± 7 ml·kg⁻¹·min⁻¹) completed preliminary tests to determine the $\dot{V}O_2$ -power output relationship and $\dot{V}O_{2\max}$. Subsequently, AOD and W' were determined from AOD, and the work completed above critical power, respectively, in CWR and 3MT. There were no differences between tests for duration, work, or average power output ($p \geq 0.05$). AOD was greatest in the CWR test (4.18 ± 0.95 vs. 3.68 ± 0.98 L; $p = 0.004$), whereas W' was greatest in the 3MT (9.55 ± 4.00 vs. 11.37 ± 3.84 kJ; $p = 0.010$). AOD and W' demonstrated a significant correlation for both CWR ($p < 0.001$, $r = 0.654$) and 3MT ($p < 0.001$, $r = 0.654$). In conclusion, despite strong correlations between AOD and W' in CWR and 3MTs, between-test differences in the magnitude of AOD and W' , suggests that the measures have different underpinning mechanisms.

Abstract word count: 197

Key words: MAOD, high-intensity, anaerobic work capacity, anaerobic

40 Introduction

41 At the onset of exercise, ATP in skeletal muscle is continuously resynthesised by the complex and
 42 closely integrated interaction of aerobic and anaerobic energy pathways (Gastin, 2001). However,
 43 whilst aerobic energy production is relatively easy to quantify as the rate of oxygen uptake at the
 44 mouth ($\dot{V}O_2$) (Poole et al., 1991), quantification of anaerobic energy production remains challenging
 45 (Noordhof, de Koning, & Foster, 2010; Noordhof, Skiba, & de Koning, 2013). Direct methods for
 46 quantifying anaerobic capacity are invasive and/or expensive, and, as a consequence, anaerobic
 47 capacity is more commonly estimated using indirect tests (Noordhof et al., 2013).

48 A common test to estimate anaerobic capacity is the accumulated oxygen deficit (AOD), as proposed
 49 by Medbø et al. (1988). The AOD determines the difference between the accumulated oxygen
 50 demand and the accumulated oxygen uptake and can be determined from a constant work-rate test to
 51 exhaustion (CWR) at a supramaximal intensity (i.e. above maximal $\dot{V}O_2$ [$\dot{V}O_{2max}$]); or an all-out test of
 52 known duration. In order to be considered as a measure of anaerobic capacity, AOD needs to reach
 53 its maximum value. Using a supramaximal CWR, it has been shown that the highest AOD is attained
 54 in tests lasting 2-5 min, which corresponds to intensities of 110-120% of $\dot{V}O_{2max}$ (Medbø et al., 1988;
 55 Muniz-Pumares, Pedlar, Godfrey, & Glaister, 2006; Weber & Schneider, 2001). The AOD, determined
 56 during all-out efforts, also appears to be sensitive to the duration of the test. All-out tests shorter than
 57 60 s tend to underestimate anaerobic capacity. Instead, if the all-out effort lasts 60-90 s, the AOD
 58 seems to plateau and reach its maximum value (Calbet, Chavarren, & Dorado, 1997; Gastin, Costill,
 59 Lawson, Krzeminski, & McConell, 1995; Withers et al., 1991; Withers, Ploeg, & Finn, 1993). The
 60 effect of all-out efforts longer than 90 s on the AOD has not been studied. It is important to note that
 61 the AOD relies on the assumptions that i) the oxygen demand can be extrapolated from the $\dot{V}O_2$ -
 62 power output relationship determined at submaximal intensities; and ii) for a given power output, the
 63 required oxygen demand is not altered during high-intensity exercise. Whilst both assumptions have
 64 been questioned, and are considered to be a limitation of the test, the AOD is considered to be the
 65 best non-invasive test to estimate anaerobic capacity (Noordhof et al., 2010).

66 Another approach to estimate anaerobic capacity has been derived from the parameters of the
 67 hyperbolic power output-duration relationship. The first component is the asymptote of the hyperbola,
 68 termed critical power, which represents the boundary between the 'heavy' and 'severe' exercise
 69 domains (Hill, 1993; Jones, Vanhatalo, Burnley, Morton, & Poole, 2010; Poole, Burnley, Vanhatalo,
 70 Rossiter, & Jones, 2016). The second component is the curvature constant (W'), which represents a
 71 fixed amount of work that can be performed above critical power (Chidnok et al., 2013; Morton, 2006).
 72 Traditionally, W' has been described as 'anaerobic work capacity', and thought to represent work
 73 produced using anaerobic energy sources (e.g. Hill, 1993; Morton, 2006). However, it has been
 74 recently suggested that the precise aetiology of W' may be more complex than originally thought,
 75 leaving its underpinning mechanisms unresolved (Broxterman et al., 2015; Dekerle et al., 2015;
 76 Murgatroyd, Ferguson, Ward, Whipp, & Rossiter, 2011; Poole et al., 2016; Simpson et al., 2015;

Skiba, Chidnok, Vanhatalo, & Jones, 2012). Nonetheless, W' is affected by glycogen content (Miura, Sato, Whipp, & Fukuba, 2000) and creatine supplementation (Smith, Stephens, Hall, Jackson, & Earnest, 1998). Moreover, W' depletion results in the build-up of fatigue-inducing metabolites associated with anaerobic energy production (Jones, Wilkerson, DiMenna, Fulford, & Poole, 2008; Poole, Ward, Gardner, & Whipp, 1988), and the rate of accumulation of those metabolites is proportional to the rate of W' depletion (Vanhatalo, Fulford, DiMenna, & Jones, 2010). As a result, the magnitude of W' typically remains constant irrespective of its rate of depletion (Chidnok et al., 2013; Fukuba et al., 2003; cf. Dekerle et al., 2015; Jones, Wilkerson, Vanhatalo, & Burnley, 2008).

The traditional method of determining W' was to model the results of 4-6 bouts of CWR exercise to exhaustion. However, the time-consuming demands of the protocol makes the approach very impractical. More recently, Vanhatalo et al. (2007) observed that the end-power output during a 3-min all-out test (3MT) corresponded to critical power; whilst the work performed above end-power output corresponded to W' . If this new approach to determining W' is valid, it should produce the same strong positive correlations with AOD as those reported when W' is determined using the traditional approach (Chatagnon, Pouilly, Thomas, & Busso, 2005; Miura, Endo, Sato, Barstow, & Fukuba, 2002),

The aims of this study, therefore, were i) to determine whether AOD and W' remain constant irrespective of their rate of depletion (i.e. CWR vs. 3MT); and ii) to investigate the relationship between AOD and W' during CWR and 3MT. It was hypothesised that both the AOD and W' would not be affected by the exercise mode. It was also hypothesised that W' and AOD would be strongly and positively correlated in both the CWR and 3MT.

Methods

Participants

Twenty-one trained male cyclists and triathletes volunteered to participate in this study, which was approved by St Mary's University Ethics Committee. Their mean \pm standard deviation (SD) for age, height and mass were 40 ± 6 years, 1.81 ± 0.08 m and 79.8 ± 7.5 kg, respectively. The participants were recruited from local cycling and triathlon clubs and can be classified as 'trained' (performance level 3; De Pauw et al., 2013). All participants provided written informed consent.

Procedures

The study consisted of four trials in an exercise physiology laboratory with controlled environmental conditions (19 ± 1 °C; $33 \pm 5\%$ relative humidity). All tests were performed on an electromagnetically braked cycle-ergometer (Lode Excalibur Sport, Groningen, Netherlands). The cycle-ergometer was individually adjusted for cyclists comfort and performance. All subsequent tests were performed using the same settings on the cycle-ergometer and at approximately the same time of the day (± 1 h). After two preliminary trials to determine the gas exchange threshold (GET), the $\dot{V}O_2$ -power output relationship, and $\dot{V}O_{2max}$, participants completed a CWR at 112.5% of $\dot{V}O_{2max}$ and a 3MT. All trials

were separated by at least 48 h to allow complete recovery. The participants were provided with a food record diary and were advised to follow a similar diet and to avoid strenuous exercise in the 24 h before each trial. Similarly, they were requested to avoid caffeine and alcohol ingestion 12 h before each trial.

Preliminary tests

The preliminary tests included two trials. In Trial 1, participants completed a ramp test to exhaustion. The test started with 3 min of unloaded cycling. The resistance of the flywheel increased thereafter at a constant rate of 30 W·min⁻¹ until exhaustion, defined in this study as a decrease in cadence of > 10 rpm for > 5 s despite strong verbal encouragement. The cadence was freely chosen by each participant and kept constant throughout the test. The preferred cadence was recorded and replicated in subsequent trials. The GET was independently identified by two investigators using the V-slope method (Beaver, Whipp, & Wasserman, 1986), and the average of the two values was used for subsequent calculations. In instances where GET estimates differed by > 10%, a third investigator determined the GET, and the average of the two closest estimates was used for analysis. Trial 2 consisted of 10 × 3-min consecutive steps to determine the relationship between $\dot{V}O_2$ and power output, followed by a ramp test to exhaustion to determine $\dot{V}O_{2max}$. The first step was performed at 50% GET and the intensity increased by 10% GET in each subsequent step, so that the final work rate corresponded to 140% GET. Steps were interspersed with 30 s of rest to allow a capillary blood sample to be drawn from the earlobe using a 20 μ L tube (EKF Diagnostics, Barleben, Germany). Whole blood samples were introduced in a pre-filled tube and analysed for blood lactate concentration (BLa) using an enzymatic-amperometric method (Biosen C-line, EKF Diagnostic, Germany). After completion of the final step, participants were allowed 5 min of stationary rest on the ergometer. Cycling was resumed at 70% GET, and increased at a rate of 15% GET every minute until volitional exhaustion (as defined above). $\dot{V}O_{2max}$ was determined as the highest $\dot{V}O_2$ obtained from a 30-s rolling average, which excluded breath-by-breath values outside 4 SD from a local (5-breath) average (Lamarra, Whipp, Ward, & Wasserman, 1987).

Constant-work rate test to exhaustion

The CWR commenced with 3 min of unloaded cycling followed by 5 min at 70% GET. Then, after 5 min stationary rest on the cycle-ergometer, participants were instructed to attain their preferred cadence after a 5-second countdown. The power output during the CWR test corresponded to 112.5% $\dot{V}O_{2max}$, determined from linear extrapolation of the relationship between $\dot{V}O_2$ and power output. The assumption of a linear $\dot{V}O_2$ -power output relationship has been challenged, though using 3-min stages, a linear relationship has been observed during for intensities up to ~95% $\dot{V}O_{2max}$, with allows estimation of supramaximal oxygen demands with 6.7% test-retest variability (Muniz-Pumares, Pedlar, Godfrey, & Glaister, 2015). Moreover, a CWR at 112.5% has been shown to elicit the greatest AOD (Muniz-Pumares et al., 2016). $\dot{V}O_2$ values to construct the $\dot{V}O_2$ -power output relationship were determined from each stage as the highest $\dot{V}O_2$ value derived from a 30-s rolling average (see above). Participants were instructed before, and encouraged throughout the test to exercise for as

long as they possibly could, but were unaware of elapsed time or expected duration. Capillary blood samples were drawn 1, 3 and 5 min after exhaustion for BLa determination.

3-min all-out test

The 3MT was performed as outlined by Vanhatalo et al. (2007). The trial commenced with 5 min cycling at 70% GET and a further 5 min resting on the cycle ergometer. Participants then completed 3 min of unloaded pedalling at their preferred cadence. In the last 10 seconds of the unloaded phase, they were instructed to increase their cadence to 110–120 rpm. At the start of the 3MT, the cycle-ergometer switched to linear mode, so that the resistance (i.e. power output) represented a function of the cadence. The alpha factor for the linear mode was determined to elicit a power output at each participant's preferred cadence corresponding to 50% of the difference between the intensity at GET and that at the end of the ramp test (i.e. 50%Δ). The subjects were instructed before the test to attain peak power (i.e. highest cadence) as soon as possible and to maintain the highest possible cadence throughout the test. Strong verbal encouragement was provided by the same investigator throughout the duration of the test. As in the CWR test, time cues were removed from the area to prevent pacing. All participants completed one familiarization trial of the 3MT that was not included in data analysis. The criteria to deem a 3MT as valid is yet to be established. Nevertheless, it has been reported that, during a 3MT: i) peak power is typically attained within the first 10 s (Vanhatalo, Doust, & Burnley, 2007); ii) peak $\dot{V}O_2$ corresponds to 97-99% $\dot{V}O_{2max}$ (Burnley, Doust, & Vanhatalo, 2006; Sperlich, Haegerle, Thissen, Mester, & Holmberg, 2011; Vanhatalo et al., 2007), although there seems to be large intrasubject variability (Sperlich et al., 2011); iii) W' is depleted to ~5% of its initial value within the first 90 s (Vanhatalo, Doust, & Burnley, 2008); and iv) end-test cadence should be within ± 10 rpm of each participant's preferred cadence, or otherwise it may affect W' (Vanhatalo et al., 2008). As in the CWR test, capillary BLa was determined 1, 3 and 5 min after the 3MT test.

Statistical analyses

The AOD was determined as the difference between the estimated oxygen demand and accumulated oxygen uptake (Medbø et al., 1988). In the CWR test, the oxygen demand was assumed to remain constant during the test (i.e. 112.5% $\dot{V}O_{2max}$), so the accumulated oxygen demand was estimated as the product of oxygen demand and the time to exhaustion (TTE). In the 3MT, raw recording of power output (6 Hz) were averaged at 1 s intervals to produce second-by-second values. The second-by-second oxygen demand was calculated from a linear projection of the $\dot{V}O_2$ -power output relationship. Subsequently, the accumulated oxygen demand was determined as the integral of second-by-second oxygen demand. Breath-by-breath $\dot{V}O_2$ data were filtered (as described above) and linearly interpolated to produce second-by-second values. The accumulated oxygen uptake was determined as the integral of second-by-second $\dot{V}O_2$. End-exercise $\dot{V}O_2$ and oxygen demand were determined in CWR and 3MT as the average $\dot{V}O_2$ and oxygen demand, respectively, in the last 10 s of the CWR and 3MT. In the 3MT, critical power was considered to be the average power output in the last 30 s of the

test. W' was determined from the 3MT (W'_{3MT}) as the integral of power output above critical power. Assuming no change in critical power (Chidnok et al., 2013), W'_{CWR} was determined as the work completed above critical power during CWR. Figure 1 outlines the protocol to determine AOD_{CWR} , AOD_{3MT} , W'_{CWR} , and W'_{3MT} . Data are presented as mean \pm SD. Using IBM SPSS 21 (IBM Corp, Armonk, NY), physiological responses to CWR and 3MT were compared using paired samples t-tests. The magnitude of the differences between CWR and 3MT were expressed as the effect size using Cohen's d , calculated as the absolute difference between means divided by the pooled SD (Cumming, 2012). Qualitative descriptors of the effect size were as follows: negligible ($d < 0.19$), small ($d = 0.20$ – 0.49), moderate ($d = 0.50$ – 0.79), or large ($d > 0.8$). Pearson product-moment correlations were determined between AOD_{3MT} and W'_{3MT} , and between AOD_{CWR} and AOD_{3MT} . In all instances, significance was accepted at $p < 0.05$.

Figure 1 near here

Results

Preliminary tests

In the ramp test, GET occurred at 188 ± 25 W and peak power output corresponded to 397 ± 46 W, so 50% Δ was 293 ± 34 W. For the 10 \times 3 min step test, the intensity at 50% GET was 94 ± 13 W and increased by 19 ± 3 W in each step, so the final intensity was 263 ± 36 W. These work rates corresponded to intensities from $41 \pm 4\%$ to $84 \pm 7\%$ $\dot{V}O_{2max}$, and raised BLa from 0.97 mmol \cdot L $^{-1}$ at the end of the first stage to 3.93 ± 1.72 mmol \cdot L $^{-1}$ for the last stage. There was a strong linear relationship between $\dot{V}O_2$ and power output for all participants ($P < 0.001$; $r = 0.995 \pm 0.004$). In the maximal test, $\dot{V}O_{2max}$ was 4.60 ± 0.61 L \cdot min $^{-1}$ (58 ± 7 mL \cdot kg $^{-1}\cdot$ min $^{-1}$).

Constant work-rate to exhaustion and 3-min all-out tests

The results from CWR and 3MT are presented in Table 1. All participants completed a valid 3MT given that: i) peak power (645 ± 127 W) was attained at the beginning of the test (6 ± 4 s); ii) peak $\dot{V}O_2$ approached $\dot{V}O_{2max}$ ($98 \pm 5\%$ $\dot{V}O_{2max}$); iii) W' was depleted to $< 15\%$ of its initial value after 90 s ($6 \pm 4\%$); and iv) the end-test cadence was within 10 rev \cdot min $^{-1}$ of the preferred cadence (4 ± 4 rev \times min $^{-1}$). Estimations of CP and W' derived from the 3MT were 316 ± 50 W ($67 \pm 8\%\Delta$) and 11.37 ± 3.84 kJ, respectively.

Table 1 near here

Estimation of anaerobic capacity from AOD and W'

There were no differences between CWR and 3MT for duration, average power output, or work completed (Table 1). However, there were differences for both estimations of anaerobic capacity between CWR and 3MT. Specifically, W'_{3MT} was greater than W'_{CWR} (small effect) whilst AOD_{CWR} was greater than AOD_{3MT} (moderate effect) (Table 1; Figure 2). In the CWR test, the estimation of anaerobic capacity, derived from AOD, was greater than that derived from W' (Table 1; mean

difference $14 \pm 10\%$; $P < 0.001$; $d = 1.17$). In contrast, there were no differences between estimations of the anaerobic energy contribution in the 3MT derived from AOD and W' (Table 1; mean difference $3 \pm 11\%$; $P = 0.175$; $d = 1.36$). AOD and W' were significantly and positively correlated in both the CWR ($r = 0.654$; $P < 0.001$) and 3MT ($r = 0.664$; $P < 0.001$).

Figure 2 near here

Discussion

The aim of the present study was to investigate AOD and W' , two parameters suggested to estimate anaerobic capacity, during a CWR and a 3MT. The main findings of the study were that i) both AOD and W' were affected by the pacing adopted and therefore different between the CWR and 3MT; ii) the differences observed between CWR and 3MT in AOD and W' followed contrasting directions such that AOD was greatest in CWR, whilst W' was greatest in 3MT; iii) there was a positive correlation between AOD and W' ; and vi) the strength of the correlation between AOD and W' was similar irrespective of pacing (i.e. CWR vs. 3MT). These results suggest that ~43% of the variance of AOD and W' is determined by a shared factor, most likely related to anaerobic energy production. However, since both estimates of anaerobic capacity were affected by pacing, and in contrasting directions, factors other than anaerobic energy production appear to influence the magnitude of AOD and/or W' .

In the present study, both AOD and W' were sensitive to pacing, as denoted by the differences between both estimates of anaerobic capacity during CWR and 3MT. However, those differences followed contrasting directions. Previous research has shown that W' remains unaffected irrespective of its rate of depletion (Chidnok et al., 2013; Fukuba et al., 2003; Vanhatalo et al., 2008); although, it has recently been shown that sudden (Dekerle et al., 2015) or progressive (Jones et al., 2008) decreases in power output might augment W' , and therefore delay exercise intolerance. In order to accept that W' remains constant irrespective of its rate of depletion, it is necessary to assume that aerobic energy production supplies power output at intensities below critical power from the onset of exercise (Jones, Vanhatalo, Burnley, Morton, & Poole, 2010; Morton, 2006), which, in turn, implies infinitely fast $\dot{V}O_2$ kinetics (Figure 1D). Despite this limitation, it is assumed that W' remains constant during a CWR test lasting > 3 min (Morton, 2006). Indeed, Chidnok et al. (2013) observed constant W' irrespective of pacing during a 3MT and a CWR of ~3.1 min. In the current study, exercise tolerance during the CWR test fell slightly short of 3 min (~2.7 min), which might not allow for a complete depletion of W' .

The AOD is thought to reach its peak value, and therefore provide an estimate of anaerobic capacity, during CWRs in which exhaustion occurs within 2-4 min (Medbø et al., 1988) or during all-outs test of at least 60 s (Gastin et al., 1995; Withers et al., 1993). In the present study, despite CWR and 3MT meeting those two conditions, AOD_{CWR} was 12% greater than AOD_{3MT}. It is possible that, given the progressive increase in $\dot{V}O_2$ and decrease in power output observed during an all-out test, $\dot{V}O_2$ at the end of the 3MT was greater than the oxygen demand, decreasing AOD_{3MT}. However, $\dot{V}O_2$ and oxygen demand at the end of the 3MT were similar (Table 1, Figure 1), and most of the AOD occurs at the

onset of all-out tests (see Figure 1). Alternatively, at the onset of the 3MT there is a higher demand of ATP turnover which can accelerate kinetics of $\dot{V}O_2$ and, possibly, reduce the AOD (Jones et al., 2008). However, studies that have examined the effects of pacing strategies during 2-6 min trials have reported that an all-out start has no effect on the AOD (Aisbett, Lerossignol, McConell, Abbiss, & Snow, 2009; Bishop, Bonetti, & Dawson, 2002). Moreover, BLa and pH, which can also be considered markers of anaerobic energy production, remain unaffected by an all-out start (Aisbett et al., 2009; Bishop et al., 2002; Chidnok et al., 2013). In contrast, the higher BLa observed in 3MT in the current study may be indicative of a greater perturbation in the muscular *milieu* during the 3MT, which in turn would affect the $\dot{V}O_2$ kinetics (e.g. Korzeniewski & Zoladz, 2015), and therefore AOD. However, whilst the increased BLa suggests higher metabolic disturbance during the 3MT, there is evidence that all-out and CWR tests result in similar intramuscular metabolic perturbation (Burnley, Vanhatalo, Fulford, & Jones, 2010). Intramuscular metabolites were not quantified in the present study, and therefore, it is difficult to account for the effect that possible differences in the metabolic *milieu* between CWR and 3MT might contribute to explain the observed difference between AOD_{3MT} and AOD_{CWR}.

Another finding of the current study was the strong correlation observed between AOD and W', which is consistent with previous research using cycle ergometry in healthy adults (Chatagnon et al., 2005; Miura et al., 2002) and children (Leclair et al., 2010). Whilst in the above studies W' was determined from several CWRs, the present study demonstrates that the relationship holds true when W' is determined from the more time-efficient 3MT. Moreover, the strength of the correlation between AOD and W' reported previously ($0.56 \leq r \leq 0.76$) compares well with the results of the present study. Overall, results suggest that, in cycling, some 34-58% of the variance of AOD and W' is underpinned by a shared mechanism, likely related to anaerobic energy production. In contrast, the same relationship has not been observed between D', the running equivalent of W', and AOD (Bosquet, Duchene, Delhors, Dupont, & Carter, 2008; Zagatto et al., 2013). Though difficult to explain, the time constant of the primary phase and the slow component of $\dot{V}O_2$ kinetics contribute to determine both W' (Murgatroyd, Ferguson, Ward, Whipp, & Rossiter, 2011) and AOD (Rossiter, 2011), and these two parameters are different between cycling and running (Hill, Halcomb, & Stevens, 2003; Pringle, Carter, Doust, & Jones, 2002). Nevertheless, the results of the present study suggest that factors other than anaerobic energy production appear to determine the magnitude of AOD and/or W', and their relationship.

During a high-intensity bout of exercise at intensities above CP, peak $\dot{V}O_2$ has been shown to correspond with $\dot{V}O_{2max}$, irrespective of the pacing strategy adopted, by some (Aisbett, Le Rossignol, & Sparrow, 2003; Aisbett et al., 2009; Bishop et al., 2002; Burnley et al., 2006; Chidnok et al., 2013; Jones et al., 2008; Simpson et al., 2015), but not all (Bailey, Vanhatalo, DiMenna, Wilkerson, & Jones, 2011; Sawyer, Morton, Womack, & Gaesser, 2012; Vanhatalo et al., 2008), studies. In the present investigation, peak $\dot{V}O_2$ during the 3MT was ~98% $\dot{V}O_{2max}$, but it only attained ~94% $\dot{V}O_{2max}$ in the CWR test, despite the intensity being ~119% of critical power. It is possible that the relatively short duration of the CWR tests combined with the possibly slower $\dot{V}O_2$ kinetics during the CWR test (see

above) resulted in a larger anaerobic energy contribution, as denoted by a greater AOD_{CWR} (Table 1). As a result, exercise might have been terminated before $\dot{V}O_{2max}$ was reached in the CWR test.

In conclusion, this is the first study to compare two approaches to estimate anaerobic capacity (AOD and W') during CWR and 3MT. Contrary to the assumption of a constant anaerobic capacity, AOD_{CWR} and W'_{3MT} were greater than AOD_{3MT} and W'_{CWR}, respectively. Nonetheless, the correlation between AOD and W' during CWR and 3MT suggests that ~43% of the magnitude of AOD and W' is determined by a shared factor, likely linked to anaerobic energy production. Moreover, the strength of the correlation between AOD and W' seems to be consistent irrespective of the type of exercise. These results suggest that anaerobic energy production is not the sole factor contributing to the magnitude of AOD and W'. Moreover, the present study suggests that factors other than anaerobic energy production contribute to AOD and W'.

References

- Aisbett, B., Lerossignol, P., McConell, G. K., Abbiss, C. R., & Snow, R. (2009). Influence of all-out and fast start on 5-min cycling time trial performance. *Medicine & Science in Sports & Exercise*, 41, 1965–71. doi.org/10.1249/MSS.0b013e3181a2aa78
- Bailey, S. J., Vanhatalo, A., DiMenna, F. J., Wilkerson, D. P., & Jones, A. M. (2011). Fast-start strategy improves $\dot{V}O_2$ kinetics and high-intensity exercise performance. *Medicine and Science in Sports and Exercise*, 43, 457–67. doi.org/10.1249/MSS.0b013e3181ef3dce
- Beaver, W., Whipp, B. J., & Wasserman, K. (1986). A new method for detecting threshold by gas exchange anaerobic. *Journal of Applied Physiology*, 60, 2020–2027.
- Bishop, D., Bonetti, D., & Dawson, B. (2002). The influence of pacing strategy on $\dot{V}O_2$ and supramaximal kayak performance. *Medicine & Science in Sports & Exercise*, 34, 1041–1047.
- Bosquet, L., Duchene, A., Delhors, P. R., Dupont, G., & Carter, H. (2008). A comparison of methods to determine maximal accumulated oxygen deficit in running. *Journal of Sports Sciences*, 26, 663–670. doi.org/10.1080/02640410701744420
- Broxterman, R. M., Ade, C. J., Craig, J. C., Wilcox, S. L., Schlup, S. J., & Barstow, T. J. (2015). Influence of blood flow occlusion on muscle oxygenation characteristics and the parameters of the power-duration relationship. *Journal of Applied Physiology*, 118, 880–9. doi.org/10.1152/jappphysiol.00875.2014
- Burnley, M., Doust, J. H., & Vanhatalo, A. (2006). A 3-min all-out test to determine peak oxygen uptake and the maximal steady state. *Medicine and Science in Sports and Exercise*, 38, 1995–2003. doi.org/10.1249/01.mss.0000232024.06114.a6
- Burnley, M., Vanhatalo, A., Fulford, J., & Jones, A. M. (2010). Similar metabolic perturbations during all-out and constant force exhaustive exercise in humans: a ⁽³¹⁾P magnetic resonance spectroscopy study. *Experimental Physiology*, 95, 798–807. doi.org/10.1113/expphysiol.2010.052688
- Calbet, J. A. L., Chavarren, J., & Dorado, C. (1997). Fractional use of anaerobic capacity during a 30- and a 45-s Wingate test. *European Journal of Applied Physiology*, 76, 308–313.
- Chatagnon, M., Pouilly, J.-P., Thomas, V., & Busso, T. (2005). Comparison between maximal power in the power-endurance relationship and maximal instantaneous power. *European Journal of Applied Physiology*, 94, 711–717. doi.org/10.1007/s00421-004-1287-y
- Cumming, G. (2012). *Understanding the new statistics*. Hove, East Sussex: Routledge.

340 Dekerle, J., de Souza, K. M., de Lucas, R. D., Guglielmo, L. G. A., Greco, C. C., & Denadai, B. S.
341 (2015). Exercise tolerance can be enhanced through a change in work rate within the severe
342 intensity domain: work above critical power is not constant. *Plos One*, 10, e0138428.
343 doi.org/10.1371/journal.pone.0138428

344 Gastin, P. B. (2001). Energy system interaction and relative contribution during maximal exercise.
345 *Sports Medicine*, 31, 725–741.

346 Gastin, P. B., Costill, D. L., Lawson, D. L., Krzeminski, K., & McConell, G. K. (1995). Accumulated
347 oxygen deficit during supramaximal all-out and constant intensity exercise. *Medicine & Science in
348 Sports & Exercise*, 27, 255–263.

349 Jones, A. M., Vanhatalo, A., Burnley, M., Morton, R., & Poole, D. C. (2010). Critical power:
350 Implications for determination of $\dot{V}O_{2max}$ and exercise tolerance. *Medicine & Science in Sports &
351 Exercise*, 42, 1876–1890.

352 Korzeniewski, B., & Zoladz, J. A. (2015). Possible mechanisms underlying slow component of $\dot{V}O_2$
353 on-kinetics in skeletal muscle. *Journal of Applied Physiology*, 118, 1240–1249.
354 doi.org/10.1152/jappphysiol.00027.2015

355 Lamarra, N., Whipp, B. J., Ward, S. a, & Wasserman, K. (1987). Effect of interbreath fluctuations on
356 characterizing exercise gas exchange kinetics. *Journal of Applied Physiology*, 62, 2003–12.

357 Leclair, E., Borel, B., Thevenet, D., Baquet, G., Mucci, P., & Berthoin, S. (2010). Assessment of child-
358 specific aerobic fitness and anaerobic capacity by the use of the power-time relationships
359 constants. *Pediatric Exercise Science*, 22, 454–66.

360 Medbø, J. I., Mohn, A. C., Tabata, I., Bahr, R., Vaage, O., & Sejersted, O. M. (1988). Anaerobic
361 capacity determined by maximal accumulated O_2 deficit. *Journal of Applied Physiology*, 64, 50–60.

362 Miura, A., Endo, M., Sato, H., Barstow, T. J., & Fukuba, Y. (2002). Relationship between the
363 curvature constant parameter of the power-duration curve and muscle cross-sectional area of the
364 thigh for cycle ergometry in humans. *European Journal of Applied Physiology*, 87, 238–244.
365 doi.org/10.1007/s00421-002-0623-3

366 Miura, A., Sato, H., Whipp, B. J., & Fukuba, Y. (2000). The effect of glycogen depletion on the
367 curvature constant parameter of the power-duration curve for cycle ergometry. *Ergonomics*, 43,
368 133–41. doi.org/10.1080/001401300184693

369 Morton, R. H. (2006). The critical power and related whole-body bioenergetic models. *European
370 Journal of Applied Physiology*, 96, 339–354. doi.org/10.1007/s00421-005-0088-2

371 Muniz-Pumares, D., Pedlar, C., Godfrey, R., Glaister, M. (2016). Accumulated oxygen deficit during
372 exercise to exhaustion determined at different supramaximal work-rates. *International Journal of
373 Sports Physiology and Performance*, in press.

374 Muniz-Pumares, D., Pedlar, C., Godfrey, R., Glaister, M. (2015). The effect of the oxygen uptake-
375 power output relationship on the prediction of supramaximal oxygen demands. *J Sports Med Phys
376 Fitness*. In press.

377 Murgatroyd, S. R., Ferguson, C., Ward, S. a, Whipp, B. J., & Rossiter, H. B. (2011). Pulmonary O_2
378 uptake kinetics as a determinant of high-intensity exercise tolerance in humans. *Journal of Applied
379 Physiology*, 110, 1598–606. doi.org/10.1152/jappphysiol.01092.2010

380 Noordhof, D. A., Skiba, P. F., & de Koning, J. J. (2013). Determining anaerobic capacity in sporting
381 activities. *International Journal of Sports Physiology and Performance*, 8, 475–82.

382 Noordhof, D. A., de Koning, J. J., & Foster, C. (2010). The maximal accumulated oxygen deficit
383 method: a valid and reliable measure of anaerobic capacity? *Sports Medicine*, 40, 285–302.
384 doi.org/10.2165/11530390-000000000-00000

385 Poole, D. C., Burnley, M., Vanhatalo, A., Rossiter, H. B., & Jones, A. M. (In press). Critical power: An
 386 important fatigue threshold in exercise physiology. *Medicine & Science in Sports & Exercise*.
 387 doi.org/10.1249/MSS.0000000000000939

388 Poole, D. C., Schaffartzik, W., Knight, D. R., Derion, T., Kennedy, B., Guy, H. J., ... Wagner, P. D.
 389 (1991). Contribution of exercising legs to the slow component of oxygen uptake kinetics in
 390 humans. *Journal of Applied Physiology*, 71, 1245–60.

391 Rossiter, H. B. (2011). Exercise: Kinetic considerations for gas exchange. *Compr Physiol*, 1, 203–44.
 392 doi.org/10.1002/cphy.c090010

393 Sawyer, B. J., Morton, R. H., Womack, C. J., & Gaesser, G. a. (2012). $\dot{V}O_{2max}$ may not be reached
 394 during exercise to exhaustion above critical power. *Medicine & Science in Sports & Exercise*, 44,
 395 1533–8. doi.org/10.1249/MSS.0b013e31824d2587

396 Simpson, L. P., Jones, A. M., Skiba, P. F., Vanhatalo, A., Wilkerson, D., Sciences, H., & Kingdom, U.
 397 (2015). Influence of hypoxia on the power-duration relationship during high-intensity exercise.
 398 *International Journal of Sports Medicine*, 36, 113–9. doi.org/10.1055/s-0034-1389943

399 Skiba, P. F., Chidnok, W., Vanhatalo, A., & Jones, A. M. (2012). Modelling the expenditure and
 400 reconstitution of work capacity above critical power. *Medicine & Science in Sports & Exercise*, 44,
 401 1526–1532. doi.org/10.1249/MSS.0b013e3182517a80

402 Smith, J. C., Stephens, D. P., Hall, E. L., Jackson, A. W., & Earnest, C. P. (1998). Effect of oral
 403 creatine ingestion on parameters of the work rate-time relationship and time to exhaustion in high-
 404 intensity cycling. *European Journal of Applied Physiology*, 77, 360–365.

405 Vanhatalo, A., Doust, J. H., & Burnley, M. (2007). Determination of critical power using a 3-min all-out
 406 cycling test. *Medicine & Science in Sports & Exercise*, 39, 548–55.
 407 doi.org/10.1249/mss.0b013e31802dd3e6

408 Vanhatalo, A., Doust, J. H., & Burnley, M. (2008). Robustness of a 3 min all-out cycling test to
 409 manipulations of power profile and cadence in humans. *Experimental Physiology*, 93, 383–390.
 410 doi.org/10.1113/expphysiol.2007.039883

411 Vanhatalo, A., Fulford, J., DiMenna, F. J., & Jones, A. M. (2010). Influence of hyperoxia on muscle
 412 metabolic responses and the power-duration relationship during severe-intensity exercise in
 413 humans: a ^{31}P magnetic resonance spectroscopy study. *Experimental Physiology*, 95, 528–540.
 414 doi.org/10.1113/expphysiol.2009.050500

415 Weber, C. L., & Schneider, D. A. (2001). Reliability of MAOD measured at 110% and 120% of peak
 416 oxygen uptake for cycling. *Medicine & Science in Sports & Exercise*, 33, 1056–1059.

417 Withers, R. T., Ploeg, G. Van Der, & Finn, J. P. (1993). Oxygen deficits during 45, 60, 75 and 90-s
 418 maximal cycling an air braked ergometer. *European Journal of Applied Physiology*, 67, 185–191.

419 Withers, R. T., Sherman, W. M., Clark, D. G., Esselbach, P. C., Nolan, S. R., Mackay, M. H., &
 420 Brinkman, M. (1991). Muscle metabolism during 30, 60 and 90 s of maximal cycling on an air-
 421 braked ergometer. *European Journal of Applied Physiology and Occupational Physiology*, 63,
 422 354–62.

423 Zagatto, A. M., Kalva-Filho, C. A., Loures, J. P., Kaminagakura, E. I., Redkva, P. E., & Papoti, M.
 424 (2013). Anaerobic running capacity determined from the critical velocity model is not significantly
 425 associated with maximal accumulated oxygen deficit in army runners. *Science & Sports*, 28, 1–7.
 426 doi.org/10.1016/j.scispo.2013.03.001.

Figures & Tables Legends

Figures.

Figure 1. Schematic representation of the methods used to determine the accumulated oxygen deficit (AOD) and W' during a 3-min all-out (3MT) test and a constant work-rate test to exhaustion. Top panels: AOD is determined as the difference between oxygen demand (dotted lines) and oxygen uptake (solid lines) during a 3MT and a CWR test (AOD_{3MT} and AOD_{CWR} ; Panels A and B, respectively). Bottom panels: W' is determined as the area between power output (solid line) and critical power (dotted line) during a 3MT and a CWR test (W'_{3MT} and W'_{CWR} ; Panels C and D, respectively).

Figure 2. Accumulated oxygen deficit and W' during constant work-rate exercise to exhaustion and a 3-min all-out test. Individual responses (dotted lines) and group means and standard deviations are shown. * denotes significantly different from the constant work-rate test ($P < 0.05$).


Table

Table 1. Physiological responses during a constant-work rate to exhaustion and a 3-min all-out test.

446


447 Figure 1.

448



449

Figure 2.



455 Table 1.

456

457

	CWR	3MT	Difference	<i>P</i> value	Cohen's <i>d</i>
Duration (s)	164 ± 46	180 ± 0	-16 ± 46	0.127	0.70
Power output (W)	376 ± 55	376 ± 55 [#]	-1 ± 23	0.882	0.01
Work (kJ)	60.85 ± 17.30	67.72 ± 9.84	-6.88 ± 15.63	0.057	0.51
W' (kJ)	9.55 ± 4.00	11.37 ± 3.84	-1.82 ± 2.93	0.010	0.46
W' (%)	20 ± 12	17 ± 6	-3 ± 9	0.116	0.37
Acc O ₂ demand (L)	14.08 ± 4.14	15.55 ± 2.14	1.48 ± 3.56	0.071	0.47
Acc O ₂ uptake (L)	9.90 ± 3.46	11.87 ± 1.48	1.97 ± 3.34	0.013	0.80
AOD (L)	4.18 ± 0.95	3.68 ± 0.98	0.50 ± 0.71	0.004	0.51
AOD (%)	31 ± 7	23 ± 5	8 ± 9	0.001	1.34
End-exercise $\dot{V}O_2$ (L·min ⁻¹)	4.29 ± 0.63	4.48 ± 0.61	-0.20 ± 0.25	0.002	0.32
End-exercise O ₂ demand (L·min ⁻¹)	5.17 ± 0.69	4.49 ± 0.61	0.68 ± 0.25	<0.001	0.55
Peak BL _a (mmol·L ⁻¹)	10.70 ± 2.57	11.77 ± 2.94	-1.07 ± 1.85	0.015	0.39
Peak HR (beats·min ⁻¹)	166 ± 11	165 ± 11	2 ± 7	0.131	0.11

[#] : average power output during the 3MT. W' (%) and AOD (%) represent the contribution of W' and AOD to the total work done and total oxygen demand, respectively.

458

459